

Test 1 - MTH 1310

Dr. Graham-Squire, Summer 2012

Name: _____

ID Number: _____

I pledge that I have neither given nor received any unauthorized assistance on this exam.

(signature)

DIRECTIONS

1. Show all of your work and use correct notation. A correct answer with insufficient work or incorrect notation will lose points.
2. Clearly indicate your answer by putting a box around it.
3. Cell phones and computers are not allowed on this test. Calculators are allowed on all parts of the test, however you should still show all of your work.
4. Give all answers in exact form, not decimal form (that is, put π instead of 3.1415, $\sqrt{2}$ instead of 1.414, etc) unless otherwise stated.
5. If you need to use the quadratic formula, it is $\frac{-b \pm \sqrt{b^2 - 4ac}}{2a}$.
6. Make sure you sign the pledge and write your ID on both pages.
7. Number of questions = 7. Total Points = 70.

1. (10 points) Two Zaks leave a house at the same time, one goes straight north and the other goes straight west. The north-going Zak travels at 9 miles per hour and the west-going Zak travels at 12 miles per hour.
 - (a) Draw a diagram to represent the situation, and let y be the distance between the two Zaks.
 - (b) Find an expression for y in terms of t , where t is time (in hours), and $t = 0$ corresponds to when the Zaks left the house.
 - (c) Find the distance between the Zaks after 2 hours.

2. (10 points) The prison population in the US has grown sharply in the past 40 years. In the 1970s, the number of inmates in prison began to exceed the capacity that the prisons were built for. The number of prisoners (in thousands) in US prisons is approximated by the function

$$N(t) = 3.5t^2 + 26.7t + 436.2 \quad (0 \leq t \leq 10)$$

where t is measured in years, with $t = 0$ corresponding to 1983. The number of inmates for which the prisons were designed is given by

$$C(t) = 24.3t + 365 \quad (0 \leq t \leq 10)$$

where $C(t)$ is also measured in thousands.

- (a) Find an expression that shows the gap between the number of prisoners and the number of inmates the prisons were built for. Simplify your expression by combining like terms.
- (b) What was the gap in 1983? In 1990?

3. (10 points) Aya wants to build a rectangular fence as the perimeter of a coop for her backyard chickens. Since she has 8 chickens, she needs to have an area of 200 ft^2 enclosed by the fence. Let x denote the length of the chicken fence and y denote the width.
- (a) Draw a diagram of the situation, and write an equation relating x and y .
 - (b) Find a function $f(x)$ giving the total amount of fencing needed to build the rectangle in terms of only the variable x .
 - (c) What is the domain of the function f ?

4. (8 points) Calculate the limits. If the limit does not exist, write DNE and explain your reasoning. Be sure to use correct notation.

(a) $\lim_{x \rightarrow (-\infty)} \frac{5x^3 - 8x^2}{7x^3 + 3x^2 + 2x}$

(b) $\lim_{x \rightarrow \infty} \frac{3x^7 + 4x - 8}{7x^3 + x}$

5. (12 points) Calculate the limits. If the limit does not exist, write DNE and explain your reasoning. Be sure to use correct notation.


(a) $\lim_{x \rightarrow 3^-} \frac{t^2 - 9}{t^3}$

(b) $\lim_{x \rightarrow 5} \frac{x^2 - 5x}{x^2 - 10x + 25}$

(c) $\lim_{x \rightarrow 4} \frac{\sqrt{x} - 2}{x - 4}$. Hint: rationalize the numerator.

6. (15 points) Sketch a graph of the following function $f(x)$, then calculate the limits and answer the question below. If the limit does not exist, write DNE and explain your reasoning.

$$f(x) = \begin{cases} 2x - 1 & \text{if } x \leq 1 \\ 4 - x & \text{if } x > 1 \end{cases}$$



(a) $\lim_{x \rightarrow 1^+} f(x) =$

(b) $\lim_{x \rightarrow 1^-} f(x) =$

(c) $\lim_{x \rightarrow 1} f(x) =$

- (d) Is the function continuous? Explain why or why not, and use the definition of continuity in your explanation.

7. (5 points) Find an equation for the line connecting the points (2,7) and (-1, 5). Write your answer in the form $y = \underline{\hspace{2cm}}$.

Extra Credit(2 points) Let $h(x) = \frac{(x^2 + 1)^3 + 4}{x^2}$. Find functions $f(x)$ and $g(x)$ so that $h(x) = f(g(x))$.