

# Calculus III, MiniTest 4 Review

Dr. Graham-Squire, Fall 2012

1. Find the curl of the vector field  $\mathbf{F}(x, y, z) = \frac{yz\mathbf{i} - xz\mathbf{j} - xy\mathbf{k}}{y^2z^2}$ . Is  $\mathbf{F}$  conservative? If so, find  $f$  such that  $\nabla f = \mathbf{F}$ .

Ans: curl  $\mathbf{F}=0$ , so  $\mathbf{F}$  is conservative.  $f(x, y, z) = \frac{x}{yz}$  is a function such that  $\nabla f = \mathbf{F}$ .

2. Evaluate the line integrals:

(a)  $\int_C (2x - y)dx + (x + 2y)dy$  where  $C$  is given by:

(i)  $C$ : one revolution counterclockwise around the circle  $x = 3 \cos t$ ,  $y = 3 \sin t$ .

Ans: Can use Green's theorem to get  $18\pi$ .

(ii)  $C$ : the line segment from  $(0,0)$  to  $(3,-3)$ .

Ans: Have to calculate the line integral directly to get 18.

(b)  $\int_C xy dx + \frac{1}{2}x^2 dy$ , where  $C$  is the boundary of the region between the graphs of  $y = x^2$  and  $y = 1$ .

Ans: Can use either the fundamental theorem of line integrals or Green's thm. to get 0.

(c)  $\int_C y dx + x dy + \frac{1}{z} dz$  where  $C$  is the curve  $\mathbf{r}(t) = \langle t, t^2 - 3t, \frac{3}{4}t + 1 \rangle$ ,  $0 \leq t \leq 4$ .

Ans: Use Fund. theorem to get  $16 + \ln 4$ .

(d)  $\int_C (x^2 - y^2) dx + 2xy dy$ , where  $C$  is given by  $x^2 + y^2 = a^2$  ( $a$  is some constant).

Ans: Use Green's theorem to get 0.

(e)  $\int_C xy ds$  where  $C$  is the line segment from  $(0,0)$  to  $(5,4)$ .

Ans: Need to use formula for line integrals (the one that involves the arc length) to get  $\frac{20\sqrt{41}}{3}$ .

3. Find an equation for the tangent plane to the paraboloid given by

$$\mathbf{r}(u, v) = u\mathbf{i} + v\mathbf{j} + (u^2 + v^2)\mathbf{k}$$

at the point  $(1,2,5)$ .

Ans:  $-2x - 4y + z = -5$

4. Evaluate the surface integral  $\iint_S z \, dS$  over the surface given by

$$\mathbf{r}(u, v) = (u + v)\mathbf{i} + (u - v)\mathbf{j} + \sin v\mathbf{k}$$

where  $0 \leq u \leq 2$  and  $0 \leq v \leq \pi$ . You may have to use Sage/Maple to evaluate the integral.

Ans:  $\int_0^\pi \int_0^2 \sin v \sqrt{2 \cos^2 v + 4} \, du \, dv = 8.623$ .